Diabetic Retinopathy/Program 3578 Session 371:A401 Half B/C



FRACTAL-BASED OSCILLATION OF MACULAR ARTERIOGENESIS AND DROPOUT DURING PROGRESSIVE DIABETIC RETINOPATHY

Patricia Parsons-Wingerter, ** Krishnan Radhakrishnan, ** Peter K. Kaiser*

*Biological Fluid Physics, NASA Glenn Research Center REB, Cleveland, OH 44135

*Department of Pathology/Cancer Center, SOM, University of New Moxics, Albuquerque, NM 87131


*Cole Eye Institute, Cleveland Clinic Foundation, Cleveland, OH 44195 "These authors contributed equally to this work.

Purpose

To examine fractal-based remodeling of macular arterial vessels with progression of diabetic retinopathy (DR).

Methods

A binary (black/white) branching pattern of arterial vessels was extracted from the macular region within retinal images obtained by 50° fluorescein angiography (FA) of eyes diagnosed with mild, moderate, or severe nonproliferative DR (NPDR) or proliferative DR (PDR). A box of 1024 by 1024 pixels centered at the fovea centralis was overlayed upon the macular region of each 2392 by 2048 binary image. One representative image of each DR stage was selected for this preliminary study. Focusing on a region around the macula, rather than studying the entire funduscopic image, considerably reduces the analysis required for diagnosis. Using VESsel GENeration Analysis (VESGEN) software, the 1024 by 1024 arterial binary pattern was mapped automatically to measure the density of total vessel length (L), as well as the fractal dimension (D) by a box-counting algorithm. VESGEN maps and quantifies vascular pattern as a function of vessel branching generation.1-7



(A and B) Artistal tress were extracted at transfer wascalar patterns from cardinating payacitie images obtained by FA using semiaternatic computer processing." (B) A box of 1004 by 1004 pixels centered on the tives centrate is everywell upon the artistal pattern (2382 by 2046 pixels). This example is the moderate NPOH image represented by its viscous executor in Fig. 2(B).

Results

For macular arterial vessels, angiogenesis oscillated strongly with vascular dropout during progression of DR. D, and L increased significantly from mild NPDR (1.28 and 0.0096 per pixel, respectively) to moderate NPDR (1.34 and 0.0130 per pixel), decreased from moderate NPDR to severe NPDR (1.28 and 0.0095 per pixel), and again increased from severe NPDR to PDR (1.30 and 0.0108 per pixel). Previously, we showed by a similar fractal analysis^a that for the combined density of macular arteries and veins, D, decreased with progression from normal to mild NPDR.




Figure 2.—Skeletonized Images of DR Progression

Figure 2.—She isolated within allege of the Progressions.

(A to D) DR proposion from mild NPDR to PDR is illustrated by the diseletorized pattern of the arterial tree, as mapped by VESGEN schware within the square loss centered on the tower centrals, as decorbed in Figure 1 VESGEN then causabled the facult disension of D) and vessel length density d), bit these representative mapper. The cross or arterial density from severe NPDR to PDR is not as great as the increase from mild to moderate NPDR, both in visible appearance and as measured by VESGEN (per Results section). This result was also obtained by the VESGEN generational branching analysis reported in Ref. 1 below.

Conclusions

By both fractal (D_i) and branching (L_i) analysis, macular arterial density oscillated with progression from mild NPDR to PDR. Results are consistent with our study reported recently for the entire arterial and venous branching trees within 50° FAs by VESGEN generational branching analysis.1 Current and previous results are important for advances in early-stage regenerative DR therapies, for which reversal of DR progression to a normal vessel density may be possible. For example, potential use of regenerative angiogenesis stimulators to reverse vascular dropout during mild and severe NPDR is not indicated for treatment of moderate NPDR.

Author Disclosure Information: P.A. Parsons-Wingerter, None; K. Radhakrishnan, None; and P.K. Kaiser, None.

References

- 1. Parsons Wingarter P, Rachaldordinans K, Volumeran M, Kalaire P, Oscillador of Angiogenesis with Valocular Diopocal in Diabetic Relencestity by VESaer GENeration Analysis (VESGEN), invest Optidisation Via Sci 2010, 51,488-507.

 2. Volumeran MB, Neth RY, McKey TL, Genicon DJ, Walanabe M, Montanas M, Kanaranari D, Kalaire PK, Seura JE, Elsevierin Q, Roba D, Hydon AD, Parsons-Wingsrisch Pr VESGEN 2D, automated, user-interaction software for quantity states and interaction. Anal Roc A 2009, 292:330-332.

 3. McKey TL, Codoon DJ, Volumeran MB, Hydon AD, Ribba D, Our HH, Kalaire PK, Parsons-Wingsrisch PF Selective in relocation of angiogenesis in small blood vessels and decrease in vessel demonstration and control for the suscular teer by transcription acceptation. Invest Capital VIII Sci 2008, 491-1184-1109.

 4. Parsons-Wingsrisch P, McKey TL, Coronie D, Volumeran MB, Condrich TK, Discovisio PE; Lymphangsgenesis by blind ended vessels sprouting is coocurrent with instrusyogenesis by viscoular spiriting. Anal Roc A 2009, 288:233-247.

 5. Patrons-Wingsrisch P, Chardonis-Maran LM, McKey TL, Rachalderham K, DiScovisio PE; Lymphangsgenesis by blind ended vessel sprouting is coocurrent with instrusyogenesis by viscoular spiriting. Anal Roc A 2009, 288:233-247.

 5. Patrons-Wingsrisch P, Chardonis-Maran LM, McKey TL, Rachalderham K, DiScovisio PE; Lymphangsgenesis by recommendation in instrusional vessel density to increased visual diameter and increased enclothile NOS activity. Microsci.

- 2004, 7:251-100.
 Place Hyportar P, Elliott KE, Clark JI, Fair AG: Fizirotiset growth factor 2 executively stimulated angiogenesis of small vessels in arterial test. Aneroscier Thromb Visio Biol 2000, 20:1250-1256.
 Passers Wingorier P, Elliott KE, Fair AG: Fizirotiset growth factor 2 executively stimulated angiyes executive in indicate state of angiogenesis in vivo by solicitor decreases in the number of new vessels. Microsise: Rus 2000, 59:221-232.
 Austral A, Rushina FE, Sogie EH, Rombris AM Clark ELIO KE, Character GE, Liciak A, Haway J-H, Parisch and rosport segon based visional roburge in the number and non-producative distinct infant. Curr Eye Res 2002, 94:274-280.